

## 6 Biomes

### 6.1 Scenarios

Since the pre-industrial simulations are an important part of the experiments, the spin-up has to finish before the pre-industrial simulations start. The spin-up should be using pre-industrial climate (**picontrol**) and year 1860 levels of “other human influences”. For this reason, the pre-industrial climate data should be replicated as often as required. The precise implementation of the spin up will be model specific, the description of which will be part of the reporting process.

Climate & CO <sub>2</sub> scenarios	
<b>picontrol</b>	Pre-industrial climate and 286ppm CO <sub>2</sub> concentration. The climate data for the entire period (1661-2299) are unique – no (or little) recycling of data has taken place.
<b>historical</b>	Historical climate and CO <sub>2</sub> concentration.
<b>rcp26</b>	Future climate and CO <sub>2</sub> concentration from RCP2.6.
<b>rcp60</b>	Future climate and CO <sub>2</sub> concentration from RCP6.0.
<b>rcp85</b>	Future climate and CO <sub>2</sub> concentration from RCP8.5.
<b>2005co2</b>	CO <sub>2</sub> concentration fixed at 2005 levels at 378.81ppm.
Human influence and land-use scenarios	
<b>1860soc</b>	Constant pre-industrial (1860) land use, nitrogen deposition, and fertilizer input.
<b>histsoc</b>	Varying historical land use, nitrogen deposition and fertilizer input.
<b>2005soc</b>	Fixed year-2005 land use, nitrogen deposition and fertilizer input.
<b>rcp26soc</b>	Varying land use, water abstraction, nitrogen deposition and fertilizer input according to SSP2 and RCP2.6.
<b>rcp60soc</b>	Varying land use, water abstraction, nitrogen deposition and fertilizer input according to SSP2 and RCP6.0.
<b>2100rcp26soc</b>	Land use, nitrogen deposition and fertilizer input fixed at year 2100 levels according to RCP2.6 in 2100.

Table 13 ISIMIP2b scenarios for the global biomes simulations.

Experiment		Input	Pre-industrial 1661-1860	Historical 1861-2005	Future 2006-2099	Extended future 2100-2299
I	no climate change, pre-industrial CO <sub>2</sub>	Climate & CO <sub>2</sub>	<b>picontrol</b>	<b>picontrol</b>	<b>picontrol</b>	<b>picontrol</b>
	varying LU & human influences up to 2005, then fixed at 2005 levels thereafter	Human & LU	<b>1860soc</b>	<b>histsoc</b>	<b>2005soc</b>	<b>2005soc</b>
II	RCP2.6 climate & CO <sub>2</sub>	Climate &	Experiment I	<b>historical</b>	<b>rcp26</b>	<b>rcp26</b>

		CO <sub>2</sub>				
	varying LU & human influences up to 2005, then fixed at 2005 levels thereafter	Human & LU		histsoc	2005soc	2005soc
IIa	RCP2.6 climate, CO <sub>2</sub> after 2005 fixed at 2005 levels	Climate & CO <sub>2</sub>	Experiment I	Experiment II	rcp26, 2005co2	rcp26, 2005co2
	varying LU & human influences up to 2005, then fixed at 2005 levels thereafter	Human & LU			2005soc	2005soc
III	RCP6.0 climate & CO <sub>2</sub>	Climate & CO <sub>2</sub>	Experiment I	Experiment II	rcp60	not simulated
	varying LU & human influences up to 2005, then fixed at 2005 levels thereafter	Human & LU			2005soc	
IV	no climate change, pre-industrial CO <sub>2</sub>	Climate & CO <sub>2</sub>	Experiment I	Experiment I	picontrol	picontrol
	varying human influences & LU up to 2100 (RCP2.6), then fixed at 2100 levels thereafter	Human & LU			rcp26soc	2100rcp26soc
V	no climate change, pre-industrial CO <sub>2</sub>	Climate & CO <sub>2</sub>	Experiment I	Experiment I	picontrol	not simulated
	varying human influences & LU (RCP6.0)	Human & LU			rcp60soc	
VI	RCP2.6 climate & CO <sub>2</sub>	Climate & CO <sub>2</sub>	Experiment I	Experiment II	rcp26	rcp26
	varying human influences & LU up to 2100 (RCP2.6), then fixed at 2100 levels thereafter	Human & LU			rcp26soc	2100rcp26soc
VII	RCP6.0 climate & CO <sub>2</sub>	Climate & CO <sub>2</sub>	Experiment I	Experiment II	rcp60	not simulated
	varying human influences & LU (RCP6.0)	Human & LU			rcp60soc	
VIII	RCP8.5 climate & CO <sub>2</sub>	Climate & CO <sub>2</sub>	Experiment I	Experiment II	rcp85	not simulated
	varying LU & human influences up to 2005, then fixed at 2005 levels thereafter	Human & LU			2005soc	

**Table 14** Additional sector-specific simulations for the biome sector.

Experiment		Input	Pre-industrial 1661-1860	Historical 1861-2005	Future 2006-2099	Extended future 2100-2299
Ia	no climate change, pre-industrial CO <sub>2</sub>	Climate & CO <sub>2</sub>	picontrol	picontrol	picontrol	picontrol
	LU & human influences fixed at 1860 levels	Human & LU	1860soc	1860soc	1860soc	1860soc
IIb	RCP2.6 climate & CO <sub>2</sub>	Climate & CO <sub>2</sub>	Experiment I	historical	rcp26	rcp26
	LU & human influences fixed at 1860 levels	Human & LU		1860soc	1860soc	1860soc
IIIa	RCP6.0 climate, CO <sub>2</sub> after 2005 fixed at 2005 levels	Climate & CO <sub>2</sub>	Experiment I	Experiment II	rcp60, 2005co2	not simulated
	varying LU & human influences up to 2005, then fixed at 2005 levels thereafter	Human & LU			2005soc	
IIIb	RCP6.0 climate & CO <sub>2</sub>	Climate & CO <sub>2</sub>	Experiment I	Experiment II	rcp60	not simulated
	LU & human influences fixed at 1860 levels	Human & LU			1860soc	
IIIc	RCP8.5 climate, CO <sub>2</sub> after 2005 fixed at 2005 levels	Climate & CO <sub>2</sub>	Experiment I	Experiment II	rcp85, 2005co2	not simulated
	varying LU & human influences up to 2005, then fixed at 2005 levels thereafter	Human & LU			2005soc	

## 6.2 Output data

**Table 15** Variables to be reported by biomes models. Variables marked by \* are also relevant for the permafrost sector and also listed in **Table 40**. **Note:** If you cannot provide the data at the temporal or spatial resolution specified, please provide it the highest possible resolution of your model.

Variable (long name)	Variable name	Unit (NetCDF name)	Resolution	Comment
<b>Essential outputs</b>				
Pools				
*Carbon Mass in Vegetation biomass	<b>cveg-&lt;pft&gt;</b>	kg m-2	per pft and gridcell total annual	Gridcell total cveg is essential. Per PFT information is desirable.
*Carbon Mass in aboveground vegetation biomass	<b>cvegag-&lt;pft&gt;</b>	kg m-2	per pft and gridcell total annual	Gridcell total cvegag is essential. Per PFT information is desirable.

*Carbon Mass in belowground vegetation biomass	<b>cvegbg-&lt;pft&gt;</b>	kg m-2	per pft and gridcell total	annual	Gridcell total cvegbg is essential. Per PFT information is desirable.
*Carbon Mass in Litter Pool	<b>clitter</b>	kg m-2	per gridcell total	annual	Info for each individual pool.
*Carbon Mass in Soil Pool	<b>csoil</b>	kg m-2	per gridcell total	annual	Info for each individual pool.
*Total Carbon Mass in Soil Pool	<b>soilc</b>	kg m-2	per gridcell total	annual	Integrated over the entire soil depth
<b>Fluxes</b>					
*Carbon Mass Flux out of atmosphere due to Gross Primary Production on Land	<b>gpp</b>	kg m-2 s-1	gridcell total	daily (monthly)	
*Carbon Mass Flux out of atmosphere due to Gross Primary Production on Land	<b>gpp-&lt;pft&gt;</b>	kg m-2 s-1	per pft	annual	
*Carbon Mass Flux into atmosphere due to Autotrophic (Plant) Respiration on Land	<b>ra</b>	kg m-2 s-1	gridcell total	daily (monthly)	
*Carbon Mass Flux out of atmosphere due to Net Primary Production on Land	<b>npp</b>	kg m-2 s-1	gridcell total	daily (monthly)	
*Carbon Mass Flux out of atmosphere due to Net Primary Production on Land	<b>npp-&lt;pft&gt;</b>	kg m-2 s-1	per pft	annual	
*Carbon Mass Flux into atmosphere due to Heterotrophic Respiration on Land	<b>rh</b>	kg m-2 s-1	gridcell total	daily (monthly)	
*Carbon Mass Flux into atmosphere due to total Carbon emissions from Fire	<b>fireint</b>	kg m-2 s-1	gridcell total	daily (monthly)	
*Carbon loss due to peat burning	<b>somcfire</b>	kg m-2 s-1	gridcell total	monthly	
*Carbon Mass Flux out of Atmosphere due to Net biome Production on Land (NBP)	<b>ecoatmflux</b>	kg m-2 s-1	gridcell total	daily (monthly)	This is the net mass flux of carbon between land and atmosphere calculated as photosynthesis MINUS the sum of plant and soil respiration, carbon fluxes from fire, harvest, grazing and land use change. Positive flux is into the land.
<b>Structure</b>					

*Leaf Area Index	<b>lai-&lt;pft&gt;</b>	1	per pft	annual	
*Leaf Area Index	<b>lai</b>	1	gridcell average	daily (monthly)	
*Plant Functional Type Grid Fraction	<b>pft-&lt;pft&gt;</b>	%	per gridcell	annual (or once if static)	The categories may differ from model to model, depending on their PFT definitions. This may include natural PFTs, anthropogenic PFTs, bare soil, lakes, urban areas, etc. Sum of all should equal the fraction of the grid-cell that is land. Value between 0 and 100.
<b>Hydrological variables</b>					
Total Evapo-Transpiration	<b>evap</b>	kg m-2 s-1	gridcell total	daily (monthly)	
Evaporation from Canopy (interception)	<b>intercept-&lt;pft&gt;</b>	kg m-2 s-1	gridcell total	daily (monthly)	The canopy evaporation+sublimation (if present in model). Provide at pft-level if available in the model
Water Evaporation from Soil	<b>esoil-&lt;pft&gt;</b>	kg m-2 s-1	per gridcell	daily (monthly)	Includes sublimation. Provide at pft-level if available in the model
Transpiration	<b>trans-&lt;pft&gt;</b>	kg m-2 s-1	per gridcell	daily (monthly)	Provide at pft-level if available in the model
*Runoff	<b>qtot</b>	kg m-2 s-1	per gridcell	daily** (monthly)	Total (surface + subsurface) runoff (qtot = qs + qsb). ** Especially for models also participating in the water sector. If daily resolution not possible, please provide monthly. If storage issues keep you from reporting daily data, please contact the ISIMIP team to discuss potential solutions.
*Soil Moisture	<b>soilmoist</b>	kg m-2	per gridcell	daily (monthly)	If possible, please provide soil moisture for all depth layers (i.e. 3D-field), and indicate depth in m. Otherwise, provide soil moisture of entire column.
Surface Runoff	<b>qs</b>	kg m-2 s-1	per gridcell	daily (monthly)	Total surface runoff leaving the land portion of the grid cell.
*Frozen soil moisture for each layer	<b>soilmoistfroz</b>	kg m-2	per gridcell	monthly	Please provide soil moisture for all depth levels and indicate depth in m.

*Snow depth	<b>snd</b>	m	per gridcell	monthly	Grid cell mean depth of snowpack.
*Snow water equivalent	<b>swe</b>	kg m-2	per gridcell	monthly	Total water mass of the snowpack (liquid or frozen), averaged over a grid cell.
*Annual maximum thaw depth	<b>thawdepth</b>	m	per gridcell	annual	Calculated from daily thaw depths. Please provide for purposes of permafrost sector.
<b>Other outputs</b>					
*Temperature of Soil	<b>tsl</b>	K	per gridcell	daily (monthly)	Temperature of each soil layer. Reported as "missing" for grid cells occupied entirely by "sea". Also needs depths in meters. Daily would be great, but otherwise monthly would work.
Burnt Area Fraction	<b>burntarea</b>	%	per gridcell	daily (monthly)	Area percentage of grid cell that has burned at any time of the given day/month/year (for daily/monthly/annual resolution)
Albedo	<b>albedo</b>	1	per gridcell	monthly	Average of pfts, snow cover, bare ground and water surfaces, range between 0-1
*N <sub>2</sub> O emissions into atmosphere	<b>n2o</b>	kg m-2 s-1	gridcell total	monthly	From land, not from industrial fossil fuel emissions and transport
*CH <sub>4</sub> emissions into atmosphere	<b>ch4</b>	kg m-2 s-1	gridcell total	monthly	From land, not from industrial fossil fuel emissions and transport

## 15 References

- Bolt, J. and van Zanden, J. L.: The Maddison Project: collaborative research on historical national accounts, *Econ. Hist. Rev.*, 67(3), 627–651, 2014.
- Choulga, M., Kourzeneva, E., Zakharova, E. and Doganovsky, A.: Estimation of the mean depth of boreal lakes for use in numerical weather prediction and climate modelling, *Tellus A Dyn. Meteorol. Oceanogr.*, 66(1), 21295, doi:10.3402/tellusa.v66.21295, 2014.
- Dellink, R., Chateau, J., Lanzi, E. and Magné, B.: Long-term economic growth projections in the Shared Socioeconomic Pathways, *Glob. Environ. Chang.*, doi:10.1016/j.gloenvcha.2015.06.004, 2015.
- Frieler, K., Lange, S., Piontek, F., Reyer, C. P. O., Schewe, J., Warszawski, L., Zhao, F., Chini, L., Denvil, S., Emanuel, K., Geiger, T., Halladay, K., Hurtt, G., Mengel, M., Murakami, D., Ostberg, S., Popp, A., Riva, R., Stevanovic, M., Suzuki, T., Volkholz, J., Burke, E., Ciais, P., Ebi, K., Eddy, T. D., Elliott, J., Galbraith, E., Gosling, S. N., Hattermann, F., Hickler, T., Hinkel, J., Hof, C., Huber, V., Jägermeyr, J., Krysanova, V., Marcé, R., Müller Schmied, H., Mouratiadou, I., Pierson, D., Tittensor, D. P., Vautard, R., van Vliet, M., Biber, M. F., Betts, R. A., Bodirsky, B. L., Deryng, D., Frohking, S., Jones, C. D., Lotze, H. K., Lotze-Campen, H., Sahajpal, R., Thonicke, K., Tian, H., and Yamagata, Y.: Assessing the impacts of 1.5 °C global warming – simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b), *Geosci. Model Dev.*, 10, 4321-4345, <https://doi.org/10.5194/gmd-10-4321-2017>, 2017.
- Haith, D. A. and Shoemaker, L. L.: Generalized Watershed Loading Functions for stream flow nutrients, *Water Resour. Bull.*, 23, 471–478, 1987.
- Klein Goldewijk, K., Beusen, A., Doelman, J., and Stehfest, E.: Anthropogenic land use estimates for the Holocene – HYDE 3.2, *Earth Syst. Sci. Data*, 9, 927-953, <https://doi.org/10.5194/essd-9-927-2017>, 2017.
- Kourzeneva, E.: External data for lake parameterization in Numerical Weather Prediction and climate modeling, *Boreal Environ. Res.*, 15(2), 165–177, 2010.
- Lamarque, J. F., Dentener, F., McConnell, J., Ro, C. U., Shaw, M., Vet, R., Bergmann, D., Cameron-Smith, P., Dalsoren, S., Doherty, R., Faluvegi, G., Ghan, S. J., Josse, B., Lee, Y. H., Mackenzie, I. a., Plummer, D., Shindell, D. T., Skeie, R. B., Stevenson, D. S., Strode, S., Zeng, G., Curran, M., Dahl-Jensen, D., Das, S., Fritzsche, D. and Nolan, M.: Multi-model mean nitrogen and sulfur deposition from the atmospheric chemistry and climate model intercomparison project (ACCMIP): Evaluation of historical and projected future changes, *Atmos. Chem. Phys.*, 13(16), 7997–8018, doi:10.5194/acp-13-7997-2013, 2013a.
- Lamarque, J. F., Shindell, D. T., Josse, B., Young, P. J., Cionni, I., Eyring, V., Bergmann, D., Cameron-Smith, P., Collins, W. J., Doherty, R., Dalsoren, S., Faluvegi, G., Folberth, G., Ghan, S. J., Horowitz, L. W., Lee, Y. H., MacKenzie, I. a., Nagashima, T., Naik, V., Plummer, D., Righi, M., Rumbold, S. T., Schulz, M., Skeie, R. B., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., Voulgarakis, a. and Zeng, G.: The atmospheric chemistry and climate model intercomparison Project (ACCMIP): Overview and description of models, simulations and climate diagnostics, *Geosci. Model Dev.*, 6(1), 179–206, doi:10.5194/gmd-6-179-2013, 2013b.
- Lange, S.: Trend-preserving bias adjustment and statistical downscaling with ISIMIP3BASD (v1.0), *Geoscientific Model Development Discussions*, doi:10.5194/gmd-2019-36, 2019.
- Lange, S.: ISIMIP3BASD. Zenodo, doi:10.5281/ZENODO.3370448, 2019b.
- De Lary, R.: Massif des Landes de Gascogne. II – ETAT DES CONNAISSANCES TECHNIQUES, Bourdeaux., 2015.
- Lehner, B. and Döll, P.: Development and validation of a global database of lakes, reservoirs and wetlands, *J. Hydrol.*, 296(1–4), 1–22, doi:10.1016/J.JHYDROL.2004.03.028, 2004.
- Millero FJ & Poisson A: International one-atmosphere equation of state of seawater. *Deep-Sea Research*, 28, 625–629, 1981.
- Monfreda, C., Ramankutty, N. and Foley, J. A.: Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000, *Glob. Biogeochem. Cycles*, 22(GB1022), doi:10.1029/2007GB002947., 2008.

- Müller Schmied, H., Adam, L., Eisner, S., Fink, G., Flörke, M., Kim, H., Oki, T., Portmann, F. T., Reinecke, R., Riedel, C., Song, Q., Zhang, J. and Döll, P.: Impact of climate forcing uncertainty and human water use on global and continental water balance components, *Proc. Int. Assoc. Hydrol. Sci.*, 93, doi:10.5194/piahs-93-1-2016, 2016.
- Murakami, D. and Yamagata, Y.: Estimation of gridded population and GDP scenarios with spatially explicit statistical downscaling, [online] Available from: <http://arxiv.org/abs/1610.09041> (Accessed 29 May 2017), 2016.
- Popp, A., Humpenöder, F., Weindl, I., Bodirsky, B. L., Bonsch, M., Lotze-Campen, H., Müller, C., Biewald, A., Rolinski, S., Stevanovic, M. and Dietrich, J. P.: Land-use protection for climate change mitigation, *Nat. Clim. Chang.*, 4(December), 2–5, doi:10.1038/nclimate2444, 2014.
- Samir, C. and Lutz, W.: The human core of the shared socioeconomic pathways: Population scenarios by age, sex and level of education for all countries to 2100, *Glob. Environ. Chang.*, doi:10.1016/j.gloenvcha.2014.06.004, 2014.
- Schneiderman, E. M., Pierson, D. C., Lounsbury, D. G. and Zion, M. S.: Modeling the hydrochemistry of the Cannonsville watershed with Generalized Watershed Loading Functions (GWLF), *J. Am. Water Resour. Assoc.*, 38, 1323–1347, 2002.
- Stevanović, M., Popp, A., Lotze-Campen, H., Dietrich, J. P., Müller, C., Bonsch, M., Schmitz, C., Bodirsky, B., Humpenöder, F. and Weindl, I.: High-end climate change impacts on agricultural welfare, *Sci. Adv.*, 2016.
- Subin, Z. M., Riley, W. J. and Mironov, D.: An improved lake model for climate simulations: Model structure, evaluation, and sensitivity analyses in CESM1, *J. Adv. Model. Earth Syst.*, 4(1), M02001, doi:10.1029/2011MS000072, 2012.
- Wada, Y., Flörke, M., Hanasaki, N., Eisner, S., Fischer, G., Tramberend, S., Satoh, Y., Van Vliet, M. T. H., Yillia, P., Ringler, C., Burek, P. and Wiberg, D.: Modeling global water use for the 21st century: The Water Futures and Solutions (WFaS) initiative and its approaches, *Geosci. Model Dev.*, 9(1), 175–222, doi:10.5194/gmd-9-175-2016, 2016.