

12 Coastal Systems

12.1 Scenarios

Climate change affects coastal systems through rising mean and extreme sea levels, causing damages through temporary flooding and losses due to permanent submergence of land. To assess these impacts, climate scenarios have to be complemented by sea-level-rise projections. While the information about thermal expansion and dynamical changes of sea level is provided by the four GCMs considered, contributions from mountain glaciers and ice sheets have to be added from other sources, which introduces a further dimension of uncertainty (see **Section 5**). The uncertainty range introduced is substantial and at least on equal footing with the climate model and scenario uncertainty (e.g., Kopp et al. 2014). To reflect this aspect, we include an additional scenario dimension in the scenario design for this sector and sample this by providing projections for the median and 5th and 95th percentiles of the contributions from ice sheets and mountain glaciers to sea-level rise. One aspect specific to the coastal-systems sector is that impacts are extremely non-linear in and sensitive to adaptation. Impacts without adaptation are 2-3 orders of magnitudes higher than those with adaptation (Hinkel et al. 2014). This leads to the circumstance that the regions with the highest infrastructure damages under the scenarios without adaptation are actually the regions least vulnerable to sea-level rise, because it is highly cost-efficient and standard practise to protect those regions against sea-level rise. Scenarios including adaptation are therefore added to the protocol to provide projections of climate change risks including adaptation potentials.

Those models that do not account for varying societal conditions (population, GDP, protection levels etc.) should keep these fixed at year 2005 levels throughout the simulations (**2005soc** scenario in Group 1 (dashed line in **Figure 1**) + **rcp26soc** or **rcp60soc** scenario in Group 2). They only need to run the first pre-industrial period of Experiment I (1661-1860). Group 3 runs only refer to models that are able to represent future changes in societal conditions.

Climate & CO ₂ scenarios	
picontrol	Pre-industrial climate (year specific for the entire period 1661-2299).
historical	Historical climate and CO ₂ concentration.
rcp26	Future climate and CO ₂ concentration from RCP2.6.
rcp60	Future climate and CO ₂ concentration from RCP6.0.
rcp85	Future climate and CO ₂ concentration from RCP8.5.
Human influence & land-use scenarios	
1860soc	Pre-industrial society and protection.
2005soc	Representation of fixed year 2005 society and protection.
ssp2soc	Varying society and protection according to SSP2.
2100ssp2soc	Representation of fixed year 2100 society and protection according to SSP2.

Table 45 ISIMIP2b scenario specification for the simulations of impacts on coastal systems.

Experiment	Input	Pre-industrial	Historical	Future	Extended future
------------	-------	----------------	------------	--------	-----------------

		1661-1860		1861-2005	2006-2099	2100-2299
I	no climate change, pre-industrial CO ₂	Climate & CO ₂	picontrol	picontrol	picontrol	picontrol
	varying society & protection up to 2005, then fixed at 2005 levels thereafter	Human & LU	Option 1: 1860soc Option 2*: 2005soc	Option 1: histsoc Option 2*: 2005soc	2005soc	2005soc
II	RCP2.6 climate & CO ₂	Climate & CO ₂	Experiment I	historical	rcp26	rcp26
	varying society & protection up to 2005, then fixed at 2005 levels thereafter	Human & LU		Option 1*: histsoc Option 2*: 2005soc	2005soc	2005soc
III	RCP6.0 climate & CO ₂	Climate & CO ₂	Experiment I	Experiment II	rcp60	not simulated
	varying society & protection up to 2005, then fixed at 2005 levels thereafter	Human & LU			2005soc	
IV	no climate change, pre-industrial CO ₂	Climate & CO ₂	Experiment I	Experiment I	picontrol	picontrol
	varying society & protection up to 2100 (SSP2), then fixed at 2100 levels thereafter	Human & LU			ssp2soc	2100ssp2soc
VI	RCP2.6 climate & CO ₂	Climate & CO ₂	Experiment I	Experiment II	rcp26	rcp26
	varying society & protection up to 2100 (SSP2), then fixed at 2100 levels thereafter	Human & LU			ssp2soc	2100ssp2soc
VII	RCP6.0 climate & CO ₂	Climate & CO ₂	Experiment I	Experiment II	rcp60	not simulated
	varying society & protection (SSP2)	Human & LU			ssp2soc	
VIII	RCP8.5 climate & CO ₂	Climate & CO ₂	Experiment I	Experiment II	rcp85	not simulated
	varying society & protection up to 2005, then fixed at 2005 levels thereafter	Human & LU			2005soc	

12.2 Output data

Table 46 Variables to be reported by coastal-systems models.

Variable (long name)	Variable name	Unit (NetCDF format)	Resolution	Comments
Expected number of people flooded annually	par	thousands/yr (1000 yr-1)	Time resolved grid	Par = People at risk.
Expected seafood costs	seafloodcost	million dollars/yr (mio 2005US\$ yr-1)		Expected annual damage caused by seafoods
Adaptation costs of building and upgrading dikes	seadikecost	million dollars/yr (mio 2005US\$ yr-1)		Cost for building/upgrading dikes
Adaptation costs of maintaining dikes	seadikemain	million dollars/yr (mio 2005US\$ yr-1)		Cost for maintenance of dikes build since the initial year (2000), but not cost for dikes “build” in the initialization of the model.

15 References

- Bolt, J. and van Zanden, J. L.: The Maddison Project: collaborative research on historical national accounts, *Econ. Hist. Rev.*, 67(3), 627–651, 2014.
- Choulga, M., Kourzeneva, E., Zakharova, E. and Doganovsky, A.: Estimation of the mean depth of boreal lakes for use in numerical weather prediction and climate modelling, *Tellus A Dyn. Meteorol. Oceanogr.*, 66(1), 21295, doi:10.3402/tellusa.v66.21295, 2014.
- Dellink, R., Chateau, J., Lanzi, E. and Magné, B.: Long-term economic growth projections in the Shared Socioeconomic Pathways, *Glob. Environ. Chang.*, doi:10.1016/j.gloenvcha.2015.06.004, 2015.¶
- Elliott, J. and Müller, C. and Deryng, D. and Chryssanthacopoulos, J. and Boote, K. J. and Büchner, M. and Foster, I. and Glotter, M. and Heinke, J. and Iizumi, T. and Izaurrealde, R. C. and Mueller, N. D. and Ray, D. K. and Rosenzweig, C. and Ruane, A. C. and Sheffield, J.: The Global Gridded Crop Model Intercomparison: data and modeling protocols for Phase 1 (v1.0), *Geosci. Model Dev.*, 8, 261–277, <https://doi.org/10.5194/gmd-8-261-2015>, 2015.
- Frieler, K., Lange, S., Piontek, F., Reyer, C. P. O., Schewe, J., Warszawski, L., Zhao, F., Chini, L., Denvil, S., Emanuel, K., Geiger, T., Halladay, K., Hurtt, G., Mengel, M., Murakami, D., Ostberg, S., Popp, A., Riva, R., Stevanovic, M., Suzuki, T., Volkholz, J., Burke, E., Ciais, P., Ebi, K., Eddy, T. D., Elliott, J., Galbraith, E., Gosling, S. N., Hattermann, F., Hickler, T., Hinkel, J., Hof, C., Huber, V., Jägermeyr, J., Krysanova, V., Marcé, R., Müller Schmied, H., Mouratiadou, I., Pierson, D., Tittensor, D. P., Vautard, R., van Vliet, M., Biber, M. F., Betts, R. A., Bodirsky, B. L., Deryng, D., Froliking, S., Jones, C. D., Lotze, H. K., Lotze-Campen, H., Sahajpal, R., Thonicke, K., Tian, H., and Yamagata, Y.: Assessing the impacts of 1.5 °C global warming – simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b), *Geosci. Model Dev.*, 10, 4321–4345, <https://doi.org/10.5194/gmd-10-4321-2017>, 2017.
- Gasparrini A, Leone M. Attributable risk from distributed lag models. *BMC Med Res Methodol.* 2014 Apr 23;14:55. doi: 10.1186/1471-2288-14-55. PMID: 24758509; PMCID: PMC4021419.¶
- Haith, D. A. and Shoemaker, L. L.: Generalized Watershed Loading Functions for stream flow nutrients, *Water Resour. Bull.*, 23, 471–478, 1987.
- Håkanson, L. Models to predict Secchi depth in small glacial lakes. *Aquatic Science* 57, 31–53 (1995). <https://doi.org/10.1007/BF00878025>¶
- Hinkel, Jochen and Lincke, Daniel and Vafeidis, Athanasios T. and Perrette, Mahé and Nicholls, Robert James and Tol, Richard S. J. and Marzeion, Ben and Fettweis, Xavier and Ionescu, Cezar and Levermann, Anders: Coastal flood damage and adaptation costs under 21st century sea-level rise, *Proceedings of the National Academy of Sciences*, 111 (9): 3292–3297; DOI: 10.1073/pnas.1222469111, 2014.¶
- Hurtt, G. C., L. Chini, R. Sahajpal, S. Froliking, B. L. Bodirsky, K. Calvin, J. C. Doelman, J. Fisk, S. Fujimori, K. K. Goldewijk, T. Hasegawa, P. Havlik, A. Heinemann, F. Humpenöder, J. Jungclaus, Jed Kaplan, J. Kennedy, T. Kristzin, D. Lawrence, P. Lawrence, L. Ma, O. Mertz, J. Pongratz, A. Popp, B. Poulter, K. Riahi, E. Shevliakova, E. Stehfest, P. Thornton, F. N. Tubiello, D. P. van Vuuren, X. Zhang (2020). Harmonization of Global Land-Use Change and Management for the Period 850–2100 (LUH2) for CMIP6. *Geoscientific Model Development Discussions*. <https://doi.org/10.5194/gmd-2019-360>
- Klein Goldewijk, K., Beusen, A., Doelman, J., and Stehfest, E.: Anthropogenic land use estimates for the Holocene – HYDE 3.2, *Earth Syst. Sci. Data*, 9, 927–953, <https://doi.org/10.5194/essd-9-927-2017>, 2017. ¶
- Kopp, Robert E. and Horton, Radley M. and Little, Christopher M. and Mitrovica, Jerry X. and Oppenheimer, Michael and Rasmussen, D. J. and Strauss, Benjamin H. and Tebaldi, Claudia: Probabilistic 21st and 22nd century sea-level projections at a global network of tide-gauge sites, *Earth's Future*, 2 (8): 383–406, <https://doi.org/10.1002/2014EF000239>, 2014.¶

- Kopp, Robert E. and Kemp, Andrew C. and Bittermann, Klaus and Horton, Benjamin P. and Donnelly, Jeffrey P. and Gehrels, W. Roland and Hay, Carling C. and Mitrovica, Jerry X. and Morrow, Eric D. and Rahmstorf, Stefan: Temperature-driven global sea-level variability in the Common Era, Proceedings of the National Academy of Sciences, 113 (11): E1434--E1441, doi:10.1073/pnas.1517056113, 2016.¶
- Kourzeneva, E. 2009. Global dataset for the parameterization of lakes in numerical weather prediction and climate modelling. ALADIN Newsletter. 37. July-December, (eds. F. Bouttier and C. Fischer), Météo-France, Toulouse, France, 46-53.
- Kourzeneva, E.: External data for lake parameterization in Numerical Weather Prediction and climate modeling, *Boreal Environ. Res.*, 15(2), 165–177, 2010.
- Lange, S.: Bias correction of surface downwelling longwave and shortwave radiation for the EWEMBI dataset, Earth Syst. Dynam., 9, 627–645, https://doi.org/10.5194/esd-9-627-2018, 2018.¶
- Lamarque, J. F., Dentener, F., McConnell, J., Ro, C. U., Shaw, M., Vet, R., Bergmann, D., Cameron-Smith, P., Dalsoren, S., Doherty, R., Faluvegi, G., Ghan, S. J., Josse, B., Lee, Y. H., Mackenzie, I. a., Plummer, D., Shindell, D. T., Skeie, R. B., Stevenson, D. S., Strode, S., Zeng, G., Curran, M., Dahl-Jensen, D., Das, S., Fritzsche, D. and Nolan, M.: Multi-model mean nitrogen and sulfur deposition from the atmospheric chemistry and climate model intercomparison project (ACCMIP): Evaluation of historical and projected future changes, *Atmos. Chem. Phys.*, 13(16), 7997–8018, doi:10.5194/acp-13-7997-2013, 2013a.
- Lamarque, J. F., Shindell, D. T., Josse, B., Young, P. J., Cionni, I., Eyring, V., Bergmann, D., Cameron-Smith, P., Collins, W. J., Doherty, R., Dalsoren, S., Faluvegi, G., Folberth, G., Ghan, S. J., Horowitz, L. W., Lee, Y. H., MacKenzie, I. a., Nagashima, T., Naik, V., Plummer, D., Righi, M., Rumbold, S. T., Schulz, M., Skeie, R. B., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., Voulgarakis, a. and Zeng, G.: The atmospheric chemistry and climate model intercomparison Project (ACCMIP): Overview and description of models, simulations and climate diagnostics, *Geosci. Model Dev.*, 6(1), 179–206, doi:10.5194/gmd-6-179-2013, 2013b.
- De Lary, R.: Massif des Landes de Gascogne. II – ETAT DES CONNAISSANCES TECHNIQUES, Bourdeaux., 2015.
- Lehner, B. and Döll, P.: Development and validation of a global database of lakes, reservoirs and wetlands, *J. Hydrol.*, 296(1–4), 1–22, doi:10.1016/J.JHYDROL.2004.03.028, 2004.
- Millero FJ & Poisson A: International one-atmosphere equation of state of seawater. *Deep-Sea Research*, 28, 625–629, 1981.
- Monfreda, C., Ramankutty, N. and Foley, J. A.: Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000, *Glob. Biogeochem. Cycles*, 22(GB1022), doi:10.1029/2007GB002947., 2008.
- Müller Schmied, H., Adam, L., Eisner, S., Fink, G., Flörke, M., Kim, H., Oki, T., Portmann, F. T., Reinecke, R., Riedel, C., Song, Q., Zhang, J. and Döll, P.: Impact of climate forcing uncertainty and human water use on global and continental water balance components, *Proc. Int. Assoc. Hydrol. Sci.*, 93, doi:10.5194/piahs-93-1-2016, 2016.
- Murakami, D. and Yamagata, Y.: Estimation of gridded population and GDP scenarios with spatially explicit statistical downscaling, [online] Available from: <http://arxiv.org/abs/1610.09041> (Accessed 29 May 2017), 2016.
- Popp, A., Humpenöder, F., Weindl, I., Bodirsky, B. L., Bonsch, M., Lotze-Campen, H., Müller, C., Biewald, A., Rolinski, S., Stevanovic, M. and Dietrich, J. P.: Land-use protection for climate change mitigation, *Nat. Clim. Chang.*, 4(December), 2–5, doi:10.1038/nclimate2444, 2014.

[Reyer, C. P. O., Silveyra Gonzalez, R., Dolos, K., Hartig, F., Hauf, Y., Noack, M., Lasch-Born, P., Rötzer, T., Pretzsch, H., Meesenburg, H., Fleck, S., Wagner, M., Bolte, A., Sanders, T. G. M., Kolari, P., Mäkelä, A., Vesala, T., Mammarella, I., Pumpanen, J., Collalti, A., Trotta, C., Matteucci, G., D'Andrea, E., Foltýnová, L., Krejza, J., Ibrom, A., Pilegaard, K., Loustau, D., Bonnefond, J.-M., Berbigier, P., Picart, D., Lafont, S., Dietze, M., Cameron, D., Vieno, M., Tian, H., Palacios-Orueta, A., Cicuendez, V., Recuero, L., Wiese, K., Büchner, M., Lange, S., Volkholz, J., Kim, H., Horemans, J. A., Bohn, F., Steinkamp, J., Chikalanov, A., Weedon, G. P., Sheffield, J., Babst, F., Vega del Valle, I., Suckow, F., Martel, S., Mahnken, M., Gutsch, M., and Frieler, K.: The PROFOUND Database for evaluating vegetation models and simulating climate impacts on European forests, *Earth Syst. Sci. Data*, 12, 1295–1320, <https://doi.org/10.5194/essd-12-1295-2020>, 2020. ¶](#)

Samir, C. and Lutz, W.: The human core of the shared socioeconomic pathways: Population scenarios by age, sex and level of education for all countries to 2100, *Glob. Environ. Chang.*, doi:10.1016/j.gloenvcha.2014.06.004, 2014.

Schneiderman, E. M., Pierson, D. C., Lounsbury, D. G. and Zion, M. S.: Modeling the hydrochemistry of the Cannonsville watershed with Generalized Watershed Loading Functions (GWLF), *J. Am. Water Resour. Assoc.*, 38, 1323–1347, 2002. ¶

[Shatwell \(unpubl.\)](#)

Stevanović, M., Popp, A., Lotze-Campen, H., Dietrich, J. P., Müller, C., Bonsch, M., Schmitz, C., Bodirsky, B., Humpenöder, F. and Weindl, I.: High-end climate change impacts on agricultural welfare, *Sci. Adv.*, 2016.

Subin, Z. M., Riley, W. J. and Mironov, D.: An improved lake model for climate simulations: Model structure, evaluation, and sensitivity analyses in CESM1, *J. Adv. Model. Earth Syst.*, 4(1), M02001, doi:10.1029/2011MS000072, 2012.

Wada, Y., Flörke, M., Hanasaki, N., Eisner, S., Fischer, G., Tramberend, S., Satoh, Y., Van Vliet, M. T. H., Yillia, P., Ringler, C., Burek, P. and Wiberg, D.: Modeling global water use for the 21st century: The Water Futures and Solutions (WFaS) initiative and its approaches, *Geosci. Model Dev.*, 9(1), 175–222, doi:10.5194/gmd-9-175-2016, 2016. ¶